## REPORT DOCUMENTATION PAGE

Form Approved

OMB NO. 0704-0188

Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing c sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any c aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations an Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-018 Washington, DC 20503.

Washington, DC 20503.		,	2	
I. AGENCY USE ONLY ( Leave Blank)	2. REPORT January 25, 1	DATE: 3. RE 2003 - Mar	PORT TYPE AND DATES COVERED: technical report, April 1, ch 31, 2003	
4. TITLE AND SUBTITLE: Evolution of Regulatory Mechanisms in Bacteria			5. FUNDING NUMBERS DAAD19-01-1-03	
6. AUTHOR(S): L. N. Ornston				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Yale University, New Haven, CT 06520		8. PEF REPO	• ^^^^	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			0\	
U. S. Army Research Office P.O. Box 12211			CY REPORT NUMBER : DAAD19-01-1-0329	
Research Triangle Park, NC 27709-2211			41868.8-25	
11. SUPPLEMENTARY NOTES				
The views, opinions and/or findings Department of the Army position, po	contained in this report olicy or decision, unless	are those of the aut so designated by o	hor(s) and should not be construed as an official ther documentation.	
12 a. DISTRIBUTION / AVAILABILITY STATEMENT			12 b. DISTRIBUTION CODE	
Approved for public release; distribution unlimited.				
that genetic analysis of structure-function rela We have extended analysis of Acinetobacter are are important environmental chemicals becau chlorogenate, a compound that resists digestic	tionships can be extended to genes to hca genes which ence se they are produced by plants on in animal feed stocks. In a	enes from any organism ode proteins that conver in response to stress. A Idition to the hca struct	ocedure for coupling PCR-mutagenesis to natural transformation as long as they provide a phenotype that is readily characterist hydroxycinnamates to protocatechuate. The hydroxxcinnama mong the chemicals degraded by hea gene products is ural genes, the Acinetobacter chromosome contains heaK which he important MarR family of transcriptional regulators that general sections in the section of the	
14. SUBJECT TERMS: bacteria, evolution, muta	orogenate, MarR 15. NUMBER OF PAGES: \$ 3			
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLAS OF ABSTRACT UNCLAS		
UNCLASSIFIED	ONCLASSIFIED		T (T	

## REPORT DOCUMENTATION PAGE (SF298) (Continuation Sheet)

- (1) List of papers submitted or published under ARO sponsorship **during this reporting period**. List the papers, including journal references, in the following categories:
  - (b) Papers published in peer-reviewed journals
- 1. **David M. Young, Ruben G. Kok, and L. Nicholas Ornston.** 2002. Phenotypic Expression of Polymerase Chain Reaction-Generated Random Mutations in a Foreign Gene After its Introduction into an *Acinetobacter* chromosome by natural transformation. Methods in Molecular Biology vol 182: In Vitro Mutagenesis Protocols, 2<sup>nd</sup> ed. J. Braman, ed.
- 2. Smith, M. A., V. B. Weaver, D. M. Young and L. N. Ornston. 2003. Genes for chlorogenate and hydroxycinnamate (*hca*) are linked to functionally related genes in the *dca-pca-qui-pob-hca* chromosomal cluster of *Acinetobacter* sp. strain ADP1. Applied and Environmental Microbiology 69:524-542.
- 2) "Scientific personnel" supported by this project and honors/awards/degrees received

Donna Parke, David Young (Ph.D. received, December 2001), Michael Smith, Nicholas Ornston

3.) Report of inventions. None

## REPORT DOCUMENTATION PAGE (SF298) (Continuation Sheet)

(4) "Scientific progress and accomplishments" (Description should include significant theoretical or experimental advances)

Acinetobacter sp. strain ADP1 is a nutritionally versatile bacterium that exhibits extraordinary competence for natural transformation. This genetic trait greatly facilitates genetic analysis and in the past has been coupled to random PCR-mutagenesis in order to identify how amino acid residues influence function in enzymes and transcriptional activators associated with catabolic pathways. We now have extended this capability to genes from other organisms, in this case Pseudomonas putida, by creating a docking site that allows PCR-amplified P. putida DNA to be integrated into the Acinetobacter chromosome under control of a promoter causing constitutive expression of genes contained in the integrated DNA. Phenotypic expression of the P. putida DNA allows identification of mutants in which the function of the P. putida gene has been altered. This development opens many opportunities for determination of how structure influences function in proteins from a full range of organisms.

Our research interest in Acinetobacter has focused upon its ability to metabolize diverse compounds that serve as nutrients for it in the natural environment. The fundamental understanding that emerges from this work is important because modification of such metabolic systems will be important in developing procedures for bioremediation. In addition, modification of the Acinetobacter pathways affords opportunities for synthesis of important chemicals by biological means.

Central to many catabolic pathways is the reaction catalyzed by protocatechuate 3,4-dioxygenase, and our past research has centered upon pcaHG, the structural genes for this enzyme. Many spontaneous mutants lacking pcaHG contain large deletions extending through flanking regions of DNA, and characterization DNA corresponding to the deletions has revealed a cluster of about 50 genes associated with dissimilation of compounds produced by plants. Among these are straight chain dicarboxylic acids and hydroxycinnamic acids which are building blocks in suberin, a protective polmer produced by plants in response to stress.

During the present research period we have focused attention on a portion of the gene cluster containing hea genes which encode proteins essential for conversion of hydroxycinnamic acids to protocatechuate. Among the enzymes involved is an unusual hydratase/lyase which is of particular interest because it converts abundant natural products such as ferulate to value added products such as the aldehyde vanillin. Among the hea genes is heaG which encodes an esterase

that cleaves chlorogenate to quinate and caffeate (caffeate is another hydroxycinnamate). Both of quinate and chlorogenate are metabolized through protocatechuate. Chlorogenate esterase is of considerable significance because its substrate presents a major problem in animal feedstocks because it is not metabolized by animals.

Downstream from the hca structural genes we discovered a gene cluster designated hcaKR. HcaK is a transporter similar in sequence to four other transporters we have discovered in our genetic analysis of Acinetobacter. HcaR is a transcriptional regulator which, unlike previous regulatory genes we have characterized, is a repressor. This is immediately significant because mutations in HcaR cause constitutive expression of enzymes that provide useful products from hydroxycinnamates. Furthermore, HcaR is a member of the MarR family of transcriptional regulators of drug resistance. We have developed physiological procedures for selection of MarR mutants, and we anticipate that analysis of such strains will provide general information about how this family of proteins exerts control over transcription.

5. Technology transfer. None.